

****FULL TITLE****

*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***

****NAMES OF EDITORS****

The Cluster-Merger Shock in 1E 0657-56: Faster than a Speeding Bullet?

Jun Koda¹, Miloš Milosavljević², Paul R. Shapiro², Daisuke Nagai³ and Ehud Nakar³

Abstract. The merging galaxy cluster 1E 0657-56, known as the “bullet cluster,” is one of the hottest clusters known. The X-ray emitting plasma exhibits bow-shock-like temperature and density jumps. The segregation of this plasma from the peaks of the mass distribution determined by gravitational lensing has been interpreted as a direct proof of collisionless dark matter. If the high shock speed inferred from the shock jump conditions equals the relative speed of the merging CDM halos, however, this merger is predicted to be such a rare event in a Λ CDM universe that observing it presents a possible conflict with the Λ CDM model.

We examined this question using high resolution, 2D simulations of gas dynamics in cluster collisions to analyze the relative motion of the clusters, the bow shock, and the contact discontinuity, and relate these to the X-ray data for the bullet cluster. We find that the velocity of the fluid shock need not equal the relative velocity of the CDM components. An illustrative simulation finds that the present relative velocity of the CDM halos is 16% lower than that of the shock. While this conclusion is sensitive to the detailed initial mass and gas density profiles of the colliding clusters, such a decrease of the inferred halo relative velocity would significantly increase the likelihood of finding 1E 0657-56 in a Λ CDM universe.

1. Introduction

We use gas dynamical simulations of the nearly head-on collision of two unequal-mass clusters to model the “bullet cluster” 1E 0657-56, to show that the high shock velocity of 4700 km s^{-1} inferred from X-ray observations (Markevitch 2006; Markevitch & Vikhlinin 2007) may exceed the relative velocity of the merging halos. Previous estimates of the extremely small probability of finding such a high-velocity cluster-cluster collision in a Λ CDM universe assumed that the shock and the halo-collision velocities were the same (Hayashi & White 2006; Farrar & Rosen 2007). Our results, first described in Milosavljević et al. (2007), significantly increase the likelihood of observing such a merger event in Λ CDM.

¹Department of Physics, University of Texas, 1 University Station C1600, Austin, TX 78712.

²Department of Astronomy, University of Texas, 1 University Station C1400, Austin, TX 78712.

³Theoretical Astrophysics, California Institute of Technology, Mail Code 130-33, 1200 East California Boulevard, Pasadena, CA 91125.

2. Simulation

We used the adaptive mesh-refinement (AMR) ASC FLASH code (Fryxell et al. 2000) in 2D cylindrical coordinates to simulate a region 20 Mpc across, with 1 kpc spatial resolution. Gravity is contributed by two rigid, spherical DM halo profiles, whose centers free-fall toward each other from rest at separation 4.6 Mpc. The density associated with the gravitational force is an NFW profile $\rho = \rho_0(r/r_s)^{-1}(1 + r/r_s)^{-2}$ for the subcluster, and a cored profile $\rho = \rho_0(1 + r/r_s)^{-3}$ for the main cluster, to imitate a non-head-on merger. The masses ($M_{500}/10^{15}M_\odot$) and radii r_{500} and r_s (Mpc) are (1.25, 1.5, 0.5) and (0.25, 0.5, 0.16) for the main cluster and subcluster, respectively.




Figure 1. Time evolution of temperature. Larger CDM halo is centered at cylindrical $(r, z) = (0, 0)$; smaller halo is at $z = 0.29, -0.66$ and -0.96 Mpc, respectively. Circles indicate the scale radii r_s of the halos.

3. Results

After subhalo “bullet” passes through the center of primary halo, the subhalo gas trails the subhalo DM, led by a bow shock and contact discontinuity (cold front), as seen in Figure 1. The opening angle and radius of curvature of the bow shock are sensitive to simulation details, but both are larger than expected for steady-state bow shocks of hard spheres with constant velocity in a uniform medium. The wings of the contact discontinuity are Kelvin-Helmholtz unstable.

During pericenter passage, the shock and contact discontinuities are slower than the relative velocity of the two CDM halos, due to ram pressure force acting on the gas. Later, the halos climb out of the gravitational potential well and decelerate, but the shock and contact discontinuity do not decelerate (Fig. 2[a]). At observed separation $D = 720$ kpc, the velocity of the shock is 4800 km s^{-1} , consistent with the 4740 km s^{-1} inferred from 1E 0657-56 data. The relative velocity of the simulated halos then is much less, 4050 km s^{-1} (16% smaller). The X-ray surface brightness and temperature profiles of the simulation viewed transverse to collision axis (Fig. 3) agree with those of 1E 0657-56.




Figure 2. (a) Velocity of the shock, V_{shock} (solid line), and of the contact discontinuity, V_{contact} (dashed line) relative to the preshock gas upstream, and the relative velocity of the CDM halos, V_{12} (dash-dot line), all as functions of the time. The distance between two CDM halos, D , is used as the coordinate of time. (b) Density and temperature jump at the shock, and the shock Mach number. (c) Position of the shock (solid line) and the contact discontinuity (dashed line) relative to subhalo center. Figures 2 and 3 are taken from Milosavljević et al. (2007).




Figure 3. (Left:) Surface brightness map of the simulated cluster in the 0.8 – 4 keV band. The circles indicate the scale radii r_s of the CDM halos. (Middle:) Temperature profile along collision axis (solid) and emissivity-weighted temperature, averaged over the (transverse) line of sight (dotted). (Right:) Surface brightness profile along collision axis.




Figure 4. Thermal bremsstrahlung emission spectrum from simulated merging cluster: the contribution from cold gas below 15 keV (dotted) and hot gas above 15 keV (short-dash), total spectrum (solid), which is the sum of cold and hot components, and single-temperature fit to cold (long-dashed) and hot (mixed dash) components. The emission from the hot shock-heated gas ($T > 15$ keV) dominates the spectrum above 20 keV and is fitted well by a single temperature profile with $T = 20$ keV.

Figure 4 shows the thermal X-ray spectrum of the simulated merging cluster. The observed spectrum of 1E 0657-56 has stronger emission above 30 keV, suggesting some additional non-thermal emission (Petrosian et al. 2006).

4. Discussion and Conclusions

Our simulations of merging galaxy clusters show that the halo collision velocity need not be the same as the intergalactic gas shock velocity. Applying our correction to the speed of the subhalo, the probability that such a massive cluster in our CDM universe has such a high-speed subhalo increases by 3 orders of magnitude relative to previous estimates, but is still small ($\sim 10^{-4}$). Springel & Farrar (2007) independently reached a similar conclusion, although their relative halo velocity is much smaller, 2600 km s^{-1} . They assumed perfectly head-on collision between a cuspy main halo and a more concentrated subhalo than ours, which would make the subhalo decelerate more before reaching the observed separation. More recent simulations by Mastropietro & Burkert (2007) show that the difference between the shock velocity and the halo relative velocity is smaller if the initial halo collision velocity is larger and if the collision is not perfectly head-on.

References

- Clowe, D., Bradač, M., Gonzalez, A. H., Markevitch, M., Randall, S. W., Jones, C., & Zaritsky, D. 2006, *ApJ*, 648, L109
 Farrar, G. R., & Rosen, R. A. 2007, *Physical Review Letters*, 98, 171302
 Fryxell, B., et al. 2000, *ApJS*, 131, 273
 Hayashi, E., & White, S. D. M. 2006, *MNRAS*, 370, L38
 Markevitch, M. 2006, *The X-ray Universe 2005*, 604, 723
 Markevitch, M., & Vikhlinin, A. 2007, *Phys. Rep.*, 443, 1
 Mastropietro, C., & Burkert, A. 2007, preprint(arXiv:0711.0967)
 Milosavljević, M., Koda, J., Nagai, D., Nakar, E., & Shapiro, P. R. 2007, *ApJ*, 661, L131
 Petrosian, V., Madejski, G., & Luli, K. 2006, *ApJ*, 652, 948
 Springel, V., & Farrar, G. R. 2007, *MNRAS*, 380, 911